

**QUIZ 2****(LESSONS 11-18: DISCRETE PROBABILITY)****MATH 119 – FALL 2023 – KUNIYUKI****100 POINTS TOTAL**

**No notes or books allowed. A scientific calculator is allowed. Simplify as appropriate. You do not have to reduce fractions. For example, 10/20 does not have to be rewritten as ½. Do not leave decimal parts in fractions or fractions in fractions, such as .1/.2 or (1/3)/(2/3).**

**MATH 119: QUIZ 2 FORMULA SHEET****Probabilities for Events Involving Equally Likely Outcomes (“elos”)**

$$P(A) = \frac{\# \text{ of elos for which } A \text{ occurs}}{N}$$

**Complementary Probabilities**

$$P(\text{not } A) = P(\bar{A}) = P(A^c) = 1 - P(A)$$

**Addition Rule for Mutually Exclusive Events (“mees”)**

$$P(A \text{ or } B) = P(A) + P(B)$$

**General Addition Rule**

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

**Multiplication Rule for Independent Events**

$$P(A \text{ and } B) = P(A) \cdot P(B)$$

$$P(A \text{ and } B \text{ and } C) = P(A) \cdot P(B) \cdot P(C)$$

**General Multiplication Rule**

$$P(A \text{ and } B) = P(A) \cdot P(B|A)$$

$$P(A \text{ and } B \text{ and } C) = P(A) \cdot P(B|A) \cdot P(C|A \text{ and } B)$$

**(SEE NEXT PAGE!)**

## Conditional Probabilities

$$P(B|A) = \frac{P(A \text{ and } B)}{P(A)}$$

$$P(B|A) = \frac{\#(A \text{ and } B)}{\#(A)} = \frac{\# \text{ of trials (or "elos") in which } A \text{ and } B \text{ occur}}{\# \text{ of trials (or "elos") in which } A \text{ occurs}}$$

## Expected Value (or Mean) of a Probability Distribution

$$E(X), \text{ or } \mu = \sum P(x) \cdot x, \text{ or } \sum x \cdot P(x)$$

## Variance and Standard Deviation of a Probability Distribution

• (NOT ON QUIZ 2)

$$VAR(X), \text{ or } \sigma^2 = \sum (x - \mu)^2 \cdot P(x), \text{ or } E(X^2) - \mu^2$$

$$SD(X), \text{ or } \sigma = \sqrt{VAR(X)}$$

## Full (or Complete) Permutations of $n$ Distinct Items

$${}_n P_n = n!$$

## (Other) Partial Permutations of $n$ Distinct Items, Taken $r$ at a Time

• (NOT ON QUIZ 2, aside from maybe  $n!$  itself)

$${}_n P_r = \frac{n!}{(n-r)!}$$

## Combinations of $n$ Distinct Items, Taken $r$ at a Time; Binomial Coefficients

$${}_n C_r, \text{ or } \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

## Binomial Probability Formula

• (NOT NEEDED ON QUIZ 2, BUT MAY HELP WITH CHECKING)

$$P(x) = {}_n C_x p^x q^{n-x}, \text{ or } \binom{n}{x} p^x q^{n-x}$$

1) (17 points). Two standard six-sided dice are rolled. One die is red; the other is green. The “number” on a die is the number of holes on the side that comes up. The “total” of the two dice is the sum of the numbers on the dice. Write your answers as **exact fractions**.

- a) What is the probability of getting a 1 on the red die **and** a 3 on the green die? Your answer will be one fraction. (2 points)

- b) What is the probability that the total is 4? Show work by using a grid as in class (you don't have to draw lines) or listing ordered pairs. (5 points)

- c) What is the probability that the number on the green die is two more than the number on the red die? Show work by using a grid as in class (you don't have to draw lines) or listing ordered pairs. (5 points)

- d) What is the probability of getting a 1 on the red die **or** a 3, a 4, or a 5 on the green die? Your answer will be one fraction. Show work by using a grid as in class (you don't have to draw lines) or listing ordered pairs. (5 points)

- 2) (5 points). Dum and Dee are candidates for president of Fredonia. 65% of adult Fredonians like Dum. 55% of adult Fredonians like Dee. 40% of adult Fredonians like both. What is the probability that a randomly selected adult Fredonian likes Dum or Dee? Write your answer as a decimal, a percent, or a fraction. Your answer will be one number (or percent), not two numbers!

- 3) (15 points). A beverage company conducts a taste test with 2000 people. Each person drinks some Regular Poke and informs the company whether the person likes or does not like Regular Poke. Each person drinks some Sugar-Free Poke and informs the company whether the person likes or does not like Sugar-Free Poke. Consider the following two-way frequency (or contingency) table. Write your answers as **exact fractions**.

		<b>Likes Regular Poke?</b>		<b>Total</b>
		<b>Yes</b>	<b>No</b>	
<b>Likes Sugar-Free Poke?</b>	<b>Yes</b>	600	300	900
	<b>No</b>	900	200	1100
<b>Total</b>		1500	500	2000

- a) What is the probability that a randomly selected person taking the taste test likes Regular Poke **or** Sugar-Free Poke? Your answer will be one fraction. (5 points)
  
- b) What is the **conditional probability** that a randomly selected person taking the taste test likes Regular Poke, **given that** the person likes Sugar-Free Poke? (5 points)
  
- c) What is the **conditional probability** that a randomly selected person taking the taste test likes Sugar-Free Poke, **given that** the person likes Regular Poke? (5 points)

4) (4 points). A fair coin can only come up “heads” or “tails,” each with the same probability. A fair coin will be flipped four times. What is the probability that the coin will come up “heads” all four times? Write your answer as an exact fraction.

5) (8 points). Three cards are randomly drawn from a standard 52-card deck **without replacement** (each drawn card is immediately, permanently removed from the deck). What is the probability that all three cards are Jacks? Write your answer as an exact fraction and also round it as a decimal to three significant figures. **Show work** by writing fractions and what you do with them!  
Hint: There are originally four Jacks in the deck of cards.

6) (4 points). A student takes three classes. Let  $X$  = the number of classes the student will pass. The student gives the following **incomplete** probability distribution for  $X$ . Find the value of  $a$  to complete the probability distribution.

Value ( $x$ )	Probability $P(x)$
0	0.100
1	0.150
2	$a$
3	0.400

7) (12 points). You and your family first pay \$500 for a life insurance policy for the year. If you die during the year, your family gets \$20,000. If you do not die during the year, your family gets nothing (beyond your presence, at least). Let  $X$  = you and your family's net monetary gain as a result of the decision to purchase the policy. Based on research, you assume that your probability of dying during the year is 0.020. Assuming this is correct, find  $E(X)$  and **interpret** it, as we did in class. (Hint: Consider many customers in a similar situation as you.) Also, fill out the table:

<b>Outcome for the year</b>	<b>Value (<math>x</math>)</b>	<b>Probability <math>P(x)</math></b>
You live.		
You die.		

8) (15 points). Showing some work or notation may help with partial credit. None of the answers is “one.” Each answer is a single number (of ways).

- a) You go to a grocery store and buy cereal, milk, and bread. The store has four brands of cereal, three brands of milk, and five brands of bread. If you only care about brands, how many ways can you buy cereal, milk, and bread at the store? (5 points)

- b) You buy ice cream at an ice cream store. The store has eight toppings and you will choose four of them for your ice cream. How many ways can you then choose the toppings? The order of the toppings does not matter. (5 points)

- c) You have seven masks, all with different colors, that you will use for a particular week. You use one mask per day during the week, and you do not reuse masks between days. If you only care about mask colors and which days you wear them on, how many ways are there for you to select your masks throughout the week? (5 points)



- 9) (5 points). A student answers all five questions on a multiple-choice math quiz. Each question has four possible options: “A,” “B,” “C,” or “D,” only one of which is correct. The student guesses randomly on all questions. The random variable is the number of questions the student gets correct. As in class, give the distribution (including the type of distribution and the values of the two parameters) that best describes the random variable.
- 10) (15 points; 5 points each). It turns out that the random variable described in 9) above has the following probability distribution table.

<b>Value (<math>x</math>)</b>	<b>Probability <math>P(x)</math></b>
0	0.237
1	0.396
2	0.264
3	0.088
4	0.015
5	0.001

Find the indicated probabilities regarding the number of questions the student gets correct. Write your answers to three decimal places. Showing work can help with partial credit; for example, rewriting using an inequality or rewriting as a sum or difference of probabilities.

- a)  $P(\text{more than } 3)$

- b)  $P(\text{at least } 1)$

- c)  $P(\text{at most } 2)$